

Appendix A

Table 4 is discussed in chapter 19. Estimated numbers in this table are to 2 significant figures. * indicates a multi-species category.

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Anchoveta(=Peruvian anchovy) (<i>Engraulis ringens</i>)	8,736,862	10-29	300,000	870,000
Japanese anchovy (<i>Engraulis japonicus</i>)	1,607,856	20-22	73,000	80,000
Sandeels(=Sandlances) nei (<i>Ammodytes</i> spp)*	530,823	10	53,000	53,000
European sprat (<i>Sprattus sprattus</i>)	652,621	17	39,000	39,000
Black and Caspian Sea sprat (<i>Clupeonella cultriventris</i>)	92,852	3-4	22,000	30,000
Capelin (<i>Mallotus villosus</i>)	1,034,615	17-50	21,000	61,000
European anchovy (<i>Engraulis encrasicolus</i>)	598,248	8-38	16,000	74,000
Araucanian herring (<i>Strangomera bentincki</i>)	427,893	27-34	13,000	16,000
Stolephorus anchovies (<i>Stolephorus</i> spp)*	269,407	3-22	12,000	87,000
European pilchard(=Sardine) (<i>Sardina pilchardus</i>)	1,049,680	69-127	8,300	15,000
Southern African anchovy (<i>Engraulis capensis</i>)	235,294	8-38	6,200	29,000
Blue whiting(=Poutassou) (<i>Micromesistius poutassou</i>)	1,864,858	135-340	5,500	14,000
Pacific anchoveta (<i>Cetengraulis mysticetus</i>)	125,698	11-24	5,200	11,000
Scads nei (<i>Decapterus</i> spp)*	1,100,340	63-231	4,800	17,000
Pacific saury (<i>Cololabis saira</i>)	378,620	83	4,600	4,600
Round sardinella (<i>Sardinella aurita</i>)	445,777	100	4,500	4,500
Pacific sandlance (<i>Ammodytes personatus</i>)	176,519	46	3,900	3,900
Sardinellas nei (<i>Sardinella</i> spp)*	821,669	7-224	3,700	120,000
Atlantic herring (<i>Clupea harengus</i>)	2,165,610	100-600	3,600	22,000
Japanese jack mackerel (<i>Trachurus japonicus</i>)	300,172	15-84	3,600	20,000
Indian scad (<i>Decapterus russelli</i>)	174,162	50	3,500	3,500
Goldstripe sardinella (<i>Sardinella gibbosa</i>)	169,019	10-55	3,100	18,000
Alaska pollock(=Walleye poll.) (<i>Theragra chalcogramma</i>)	2,903,353	227-1,000	2,900	13,000
Anchovies, etc. nei (<i>Engraulidae</i>)*	268,825	3-94	2,800	87,000
California pilchard (<i>Sardinops sagax</i>)	516,144	120-183	2,800	4,300
Indian oil sardine (<i>Sardinella longiceps</i>)	370,242	87-132	2,800	4,300

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Yellowstripe scad (<i>Selaroides leptolepis</i>)	176,757	64-66	2,700	2,800
Bombay-duck (<i>Harpadon nehereus</i>)	181,818	5-69	2,600	38,000
Norway pout (<i>Trisopterus esmarkii</i>)	70,670	20-28	2,500	3,500
Chub mackerel (<i>Scomber japonicus</i>)	1,781,553	100-750	2,400	18,000
Gulf menhaden (<i>Brevoortia patronus</i>)	513,153	234	2,200	2,200
Elongate ilisha (<i>Ilisha elongata</i>)	87,018	31-42	2,100	2,800
Largehead hairtail (<i>Trichiurus lepturus</i>)	1,298,327	32-713	1,800	41,000
Chilean jack mackerel (<i>Trachurus murphyi</i>)	1,778,803	200-1,000	1,800	8,900
Japanese pilchard (<i>Sardinops sagax</i>)	263,117	120-183	1,400	2,200
Atlantic mackerel (<i>Scomber scombrus</i>)	650,054	389-454	1,400	1,700
Atlantic horse mackerel (<i>Trachurus trachurus</i>)	228,074	160	1,400	1,400
Southern African pilchard (<i>Sardinops sagax</i>)	243,062	120-183	1,300	2,000
Rainbow sardine (<i>Dussumieria acuta</i>)	35,592	23-28	1,300	1,600
Indian mackerel (<i>Rastrelliger kanagurta</i>)	226,216	54-193	1,200	4,200
Kelee shad (<i>Hilsa kelee</i>)	87,178	40-73	1,200	2,200
Whitehead's round herring (<i>Etrumeus whiteheadi</i>)	49,414	29-40	1,200	1,700
Cape horse mackerel (<i>Trachurus capensis</i>)	361,017	291	1,200	1,200
Bali sardinella (<i>Sardinella lemuru</i>)	120,128	49-112	1,100	2,400
Californian anchovy (<i>Engraulis mordax</i>)	24,102	22	1,100	1,100
Pacific thread herring (<i>Opisthonema libertate</i>)	111,174	33-110	1,000	3,400
Argentine anchovy (<i>Engraulis anchoita</i>)	27,179	4-31	890	7,600
Pacific herring (<i>Clupea pallasii pallasii</i>)	351,598	64-400	880	5,500
Madeiran sardinella (<i>Sardinella maderensis</i>)	145,688	30-224	650	4,900
Japanese scad (<i>Decapterus maruadsi</i>)	41,089	63	650	650
Silver pomfrets nei (<i>Pampus</i> spp)*	320,531	455-500	640	700
Red-eye round herring (<i>Etrumeus teres</i>)	39,976	63	640	640
Argentine hake (<i>Merluccius hubbsi</i>)	375,198	400-600	630	940
Bonga shad (<i>Ethmalosa fimbriata</i>)	181,398	143-292	620	1,300
Atlantic menhaden (<i>Brevoortia tyrannus</i>)	211,127	162-349	600	1,300
Bigeye grunt (<i>Brachydeuterus auritus</i>)	26,586	10-45	590	2,600
South American pilchard (<i>Sardinops sagax</i>)	108,655	120-183	590	910
Bigeye scad (<i>Selar crumenophthalmus</i>)	133,386	230	580	580
Yellow croaker (<i>Larimichthys polyactis</i>)	272,569	229-499	550	1,200
Falkland sprat (<i>Sprattus fuegensis</i>)	10,043	16-19	520	630
Brazilian sardinella (<i>Sardinella janeiro</i>)	37,329	72	520	520

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Pacific menhaden (<i>Ethmidium maculatum</i>)	23,980	39-53	450	610
Beaked redfish (<i>Sebastes mentella</i>)	76,115	195	390	390
Yellowfin sole (<i>Limanda aspera</i>)	73,558	200	370	370
Goatfishes (<i>Upeneus</i> spp)*	66,693	7-189	350	9,500
Jack and horse mackerels nei (<i>Trachurus</i> spp)*	340,018	11-1,000	340	30,000
Southern blue whiting (<i>Micromesistius australis</i>)	135,050	400	340	340
Whiting (<i>Merlangius merlangus</i>)	54,774	122-165	330	450
North Pacific hake (<i>Merluccius productus</i>)	278,506	500-1,035	270	560
Bogue (Boops boops)	30,319	58-112	270	520
Frigate and bullet tunas (<i>Auxis thazard</i> , <i>A. rochei</i>)*	255,070	237-1,034	250	1,100
Polar cod (<i>Boreogadus saida</i>)	27,374	83-112	250	330
Brazilian flathead (<i>Percophis brasiliensis</i>)	7,953	4-33	240	1,900
Atlantic cod (<i>Gadus morhua</i>)	898,998	800-4,000	220	1,100
Skipjack tuna (<i>Katsuwonus pelamis</i>)	2,161,174	2,000-10,000	220	1,100
Atlantic thread herring (<i>Opisthonema oglinum</i>)	18,992	65-91	210	290
Bigeyes nei (<i>Priacanthus</i> spp)*	121,170	50-618	200	2,400
Klunzinger's mullet (<i>Liza klunzingeri</i>)	5,429	20-28	200	270
Japanese sardinella (<i>Sardinella zunasi</i>)	1,998	10	190	190
Hilsa shad (<i>Tenulosa ilisha</i>)	246,823	1,400	180	180
Snakeskin gourami (<i>Trichogaster pectoralis</i>)	26,137	154	170	170
Haddock (<i>Melanogrammus aeglefinus</i>)	282,227	900-1,800	160	310
Pink(=Humpback)salmon (<i>Oncorhynchus gorboscha</i>)	357,492	1,364-2,273	160	260
Okhotsk atka mackerel (<i>Pleurogrammus azonus</i>)	208,605	1,295	160	160
Cape hakes (<i>Merluccius capensis</i> , <i>M. paradox.</i>)*	305,500	465-2,030	150	660
Cunene horse mackerel (<i>Trachurus trecae</i>)	66,436	21-498	130	3,200
Silver-stripe round herring (<i>Spratelloides gracilis</i>)	503	3-4	130	160
Blue grenadier (<i>Macruronus novaezelandiae</i>)	198,697	1,500	130	130
Mullets nei (<i>Mugilidae</i>)*	244,721	20-2,000	120	12,000
Patagonian grenadier (<i>Macruronus magellanicus</i>)	252,763	993-2,103	120	250
Nile tilapia (<i>Oreochromis niloticus niloticus</i>)	234,894	2,000	120	120
Atlantic bumper (<i>Chloroscombrus chrysurus</i>)	16,007	21-140	110	760
Butterfishes, pomfrets nei (<i>Stromateidae</i>)*	56,937	80-500	110	710
Big-scale sand smelt (<i>Atherina boyeri</i>)	963	3-9	110	290

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Black pomfret (<i>Parastromateus niger</i>)	64,381	583	110	110
Torpedo scad (<i>Megalaspis cordyla</i>)	86,578	931	93	93
Atlantic silverside (<i>Menidia menidia</i>)	424	5	92	92
Common dab (<i>Limanda limanda</i>)	16,287	166-178	91	98
Flathead grey mullet (<i>Mugil cephalus</i>)	122,175	909-1,364	90	130
Saithe(=Pollock) (<i>Pollachius virens</i>)	396,027	2,200-4,500	88	180
European plaice (<i>Pleuronectes platessa</i>)	94,826	1,100	86	86
Silver pomfret (<i>Pampus argenteus</i>)	42,526	455-500	85	93
Goatfishes, red mullets nei (<i>Mullidae</i>)*	25,723	7-313	82	3,700
Atlantic redfishes nei (<i>Sebastes</i> spp)*	104,973	109-1,284	82	960
Chacunda gizzard shad (<i>Anodontostoma chacunda</i>)	6,440	78-79	82	82
Nile perch (<i>Lates niloticus</i>)	318,445	2,000-4,000	80	160
Cabinza grunt (<i>Isacia conceptionis</i>)	4,396	40-55	80	110
Whitemouth croaker (<i>Micropogonias furnieri</i>)	80,366	638-1,023	79	130
Pacific cod (<i>Gadus macrocephalus</i>)	354,889	2,273-4,545	78	160
Bobo croaker (<i>Pseudotolithus elongatus</i>)	16,150	216	75	75
Wolf-herrings nei (<i>Chirocentrus</i> spp)*	54,007	90-757	71	600
Arrow-tooth flounder (<i>Atheresthes stomias</i>)	18,062	131-260	69	140
Atka mackerel (<i>Pleurogrammus monopterygius</i>)	51,226	697-748	68	73
Yellowfin tuna (<i>Thunnus albacares</i>)	1,257,110	5,000-20,000	63	250
Indo-Pacific king mackerel (<i>Scomberomorus guttatus</i>)	43,983	730	60	60
Surmullet (<i>Mullus surmuletus</i>)	15,153	81-260	58	190
So-iny (redlip) mullet (<i>Chelon haematocheilus</i>)	70,635	1,238	57	57
Roundnose grenadier (<i>Coryphaenoides rupestris</i>)	27,617	372-507	54	74
South Pacific hake (<i>Merluccius gayi gayi</i>)	134,328	886-2,699	50	150
European flounder (<i>Platichthys flesus</i>)	18,006	239-367	49	75
Kawakawa (<i>Euthynnus affinis</i>)	233,558	5,000	47	47
Brazilian codling (<i>Urophycis brasiliensis</i>)	5,604	118-121	46	47
Scaled sardines (<i>Harengula</i> spp)*	1,404	19-32	44	74
Common sole (<i>Solea solea</i>)	41,471	23-971	43	1,800
Whitefin wolf-herring (<i>Chirocentrus nudus</i>)	5,136	90-123	42	57
European hake (<i>Merluccius merluccius</i>)	72,895	1,800	40	40
Frigate tuna (<i>Auxis thazard thazard</i>)	39,814	237-1,034	39	170
Chum(=Keta=Dog)salmon (<i>Oncorhynchus keta</i>)	316,054	3,182-8,182	39	99

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Crucian carp (<i>Carassius carassius</i>)	5,648	150	38	38
Mediterranean horse mackerel (<i>Trachurus mediterraneus</i>)	19,823	11-600	33	1,800
Narrow-barred Spanish mackerel (<i>Scomberomorus commerson</i>)	197,585	5,750-5,900	33	34
Tadpole codling (<i>Salilota australis</i>)	10,247	210-329	31	49
Argentines (<i>Argentina spp</i>)*	31,780	56-1,079	29	560
Atlantic saury (<i>Scomberesox saurus saurus</i>)	3,158	80-110	29	39
Bleak (<i>Alburnus alburnus</i>)	800	23-28	29	35
Roach (<i>Rutilus rutilus</i>)	7,044	69-255	28	100
Yellowtail flounder (<i>Limanda ferruginea</i>)	16,521	500-600	28	33
Blue tilapia (<i>Oreochromis aureus</i>)	2,568	95	27	27
Sockeye(=Red)salmon (<i>Oncorhynchus nerka</i>)	131,284	2,300-5,000	26	57
Daggertooth pike conger (<i>Muraenesox cinereus</i>)	263,236	10,689	25	25
Black scabbardfish (<i>Aphanopus carbo</i>)	11,901	499	24	24
Japanese Spanish mackerel (<i>Scomberomorus niphonius</i>)	49,813	1,562-2,130	23	32
Alewife (<i>Alosa pseudoharengus</i>)	6,372	227-273	23	28
Argentine (<i>Argentina sphyraena</i>)	1,505	56-64	23	27
Bigeye tuna (<i>Thunnus obesus</i>)	439,009	15,000-20,000	22	29
Large yellow croaker (<i>Larimichthys crocea</i>)	73,610	451-3,490	21	160
Indian halibut (<i>Psettodes erumei</i>)	19,801	941	21	21
So-iuy mullet (<i>Mugil soiuy</i>)	4,947	67-247	20	74
Argentine croaker (<i>Umbrina canosai</i>)	14,510	394-739	20	37
Blackbelly rosefish (<i>Helicolenus dactylopterus dactylopterus</i>)	5,782	218-284	20	27
European perch (<i>Perca fluviatilis</i>)	23,493	1,200	20	20
Mozambique tilapia (<i>Oreochromis mossambicus</i>)	19,884	1,000	20	20
Pacific ocean perch (<i>Sebastes alutus</i>)	28,472	1,400	20	20
Dorab wolf-herring (<i>Chirocentrus dorab</i>)	14,605	717-757	19	20
Snoek (<i>Thysites atun</i>)	43,106	2,289	19	19
Japanese threadfin bream (<i>Nemipterus japonicus</i>)	5,500	42-309	18	130
Orange roughy (<i>Hoplostethus atlanticus</i>)	26,490	1,500	18	18
Rainbow smelt (<i>Osmerus mordax mordax</i>)	1,562	85	18	18
Witch flounder (<i>Glyptocephalus cynoglossus</i>)	16,946	300-1,000	17	56
Ballyhoo halfbeak (<i>Hemiramphus brasiliensis</i>)	2,125	134	16	16
Greenback horse mackerel (<i>Trachurus declivis</i>)	13,401	850	16	16

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Surmullets(=Red mullets) nei (Mullus spp)*	14,963	81-1,000	15	190
Silver hake (Merluccius bilinearis)	23,169	1,298-1,524	15	18
Golden threadfin bream (Nemipterus virgatus)	3,569	208-231	15	17
Pike-perch (Sander lucioperca)	16,989	146-1,200	14	120
Lemon sole (Microstomus kitt)	12,639	649-885	14	19
Freshwater bream (Abramis brama)	49,501	3,600	14	14
Cape bonnetmouth (Emmelichthys nitidus nitidus)	4,168	66-324	13	63
Atlantic croaker (Micropogonias undulatus)	11,442	227-909	13	50
Brazilian menhaden (Brevoortia aurea)	1,222	67-91	13	18
African sicklefish (Drepane africana)	2,873	165-225	13	17
Boe drum (Pteroscion peli)	1,522	98-114	13	16
Blue mackerel (Scomber australasicus)	13,012	1,000	13	13
Bluefish (Pomatomus saltatrix)	26,158	1,943	13	13
Red porgy (Pagrus pagrus)	8,366	660	13	13
Red bigeye (Priacanthus macracanthus)	2,456	50-200	12	49
Albacore (Thunnus alalunga)	233,056	4,540-21,364	11	51
Atlantic anchoveta (Cetengraulis edentulus)	278	16-25	11	18
Longtail tuna (Thunnus tonggol)	208,271	15,000-20,000	10	14
Atlantic moonfish (Selene setapinnis)	3,440	333	10	10
Bastard halibuts nei (Paralichthys spp)*	7,173	368-754	9.5	19
Barracudas nei (Sphyræna spp)*	85,722	187-9,072	9.4	460
Grey gurnard (Eutrigla gurnardus)	5,416	538-574	9.4	10
American yellow perch (Perca flavescens)	4,071	114-454	9.0	36
Atlantic pomfret (Brama brama)	10,910	737-1,260	8.7	15
Bagrid catfish (Chrysichthys nigrodigitatus)	11,272	951-1,296	8.7	12
Red mullet (Mullus barbatus barbatus)	8,603	500-1,000	8.6	17
Bigeye croaker (Pennahia anea)	1,480	172	8.6	8.6
Atlantic searobins (Prionotus spp)*	4,456	121-527	8.5	37
Greater weever (Trachinus draco)	854	78-110	7.7	11
Tigertooth croaker (Otolithes ruber)	6,788	953	7.1	7.1
Atlantic bonito (Sarda sarda)	34,255	1,818-5,000	6.9	19
John dory (Zeus faber)	9,717	1,400	6.9	6.9
Japanese eel (Anguilla japonica)	584	86	6.8	6.8
Bluefin gurnard (Chelidonichthys kumu)	3,986	603	6.6	6.6
Black seabream (Spondyliosoma cantharus)	6,424	1,000	6.4	6.4

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Burbot (<i>Lota lota</i>)	3,167	497	6.4	6.4
Goldfish (<i>Carassius auratus auratus</i>)	3,580	7-567	6.3	500
Bastard grunt (<i>Pomadasyd incisus</i>)	2,417	27-419	5.8	89
Australian pilchard (<i>Sardinops sagax</i>)	1,059	120-183	5.8	8.8
Lebranche mullet (<i>Mugil liza</i>)	3,109	550	5.7	5.7
Sablefish (<i>Anoplopoma fimbria</i>)	25,657	4,536	5.7	5.7
Tub gurnard (<i>Chelidonichthys lucernus</i>)	2,585	220-464	5.6	12
Northern pike (<i>Esox lucius</i>)	24,541	4,500	5.5	5.5
Silver scabbardfish (<i>Lepidopus caudatus</i>)	12,490	1,000-2,300	5.4	12
Amer. plaice(=Long rough dab) (<i>Hippoglossoides platessoides</i>)	13,127	909-2,500	5.3	14
Barramundi(=Giant seaperch) (<i>Lates calcarifer</i>)	73,485	1,500-14,000	5.2	49
Picked dogfish (<i>Squalus acanthias</i>)	23,160	3,000-4,500	5.1	7.7
Atlantic wolffish (<i>Anarhichas lupus</i>)	33,220	6,804	4.9	4.9
Common carp (<i>Cyprinus carpio carpio</i>)	71,339	2,000-15,000	4.8	36
Garfish (<i>Belone belone</i>)	2,335	20-500	4.7	120
African moonfish (<i>Selene dorsalis</i>)	2,328	142-494	4.7	16
North African catfish (<i>Clarias gariepinus</i>)	39,147	5,411-8,249	4.7	7.2
Pacific sierra (<i>Scomberomorus sierra</i>)	8,581	1,814	4.7	4.7
Winter flounder (<i>Pseudopleuronectes americanus</i>)	6,602	1,400	4.7	4.7
Walleye (<i>Sander vitreus</i>)	8,264	907-1,814	4.6	9.1
Greenland halibut (<i>Reinhardtius hippoglossoides</i>)	111,415	2,273-25,000	4.5	49
Angolan dentex (<i>Dentex angolensis</i>)	1,697	101-386	4.4	17
Blackmouth croaker (<i>Atroubucca nibe</i>)	412	93	4.4	4.4
Cassava croaker (<i>Pseudotolithus senegalensis</i>)	4,407	1,000	4.4	4.4
Scup (<i>Stenotomus chrysops</i>)	3,280	750	4.4	4.4
Blackbanded trevally (<i>Seriolina nigrofasciata</i>)	5,941	1,500	4.0	4.0
Benguela hake (<i>Merluccius polli</i>)	1,758	55-467	3.8	32
Rough scad (<i>Trachurus lathami</i>)	960	251	3.8	3.8
Amberjacks nei (<i>Seriola spp</i>)*	100,700	1,057-27,300	3.7	95
Bonytongues nei (<i>Heterotis spp</i>)*	11,054	2,244-3,060	3.6	4.9
Serra Spanish mackerel (<i>Scomberomorus brasiliensis</i>)	6,831	1,476-2,013	3.4	4.6
American angler (<i>Lophius americanus</i>)	22,812	3,182-6,818	3.3	7.2
Patagonian toothfish (<i>Dissostichus eleginoides</i>)	32,503	9,000-10,000	3.3	3.6

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Black cusk-eel (<i>Genypterus maculatus</i>)	1,729	523	3.3	3.3
Tusk(=Cusk) (<i>Brosme brosme</i>)	26,487	8,000	3.3	3.3
American gizzard shad (<i>Dorosoma cepedianum</i>)	1,683	18-520	3.2	93
Yellowtail snapper (<i>Ocyurus chrysurus</i>)	6,487	750-2,000	3.2	8.6
Coho(=Silver)salmon (<i>Oncorhynchus kisutch</i>)	18,889	6,000	3.1	3.1
Milkfish (<i>Chanos chanos</i>)	1,857	600	3.1	3.1
Bartail flathead (<i>Platycephalus indicus</i>)	2,894	166-969	3.0	17
Summer flounder (<i>Paralichthys dentatus</i>)	6,887	907-2,268	3.0	7.6
Atlantic Spanish mackerel (<i>Scomberomorus maculatus</i>)	7,972	907-2,722	2.9	8.8
Red codling (<i>Pseudophycis bachus</i>)	7,174	1,500-2,500	2.9	4.8
Axillary seabream (<i>Pagellus acarne</i>)	1,273	13-447	2.8	95
Alfonsinos nei (<i>Beryx spp</i>)*	8,933	596-3,239	2.8	15
African lungfishes (<i>Protopterus spp</i>)*	13,692	217-5,100	2.7	63
Greater Argentine (<i>Argentina silus</i>)	2,893	300-1,079	2.7	9.6
Little tunny(=Atl.black skipj) (<i>Euthynnus alletteratus</i>)	14,774	3,723-5,548	2.7	4.0
False scad (<i>Caranx rhonchus</i>)	2,387	34-933	2.6	69
Crevalle jack (<i>Caranx hippos</i>)	5,952	1,361-2,268	2.6	4.4
Redfish (<i>Centroberyx affinis</i>)	1,477	160-600	2.5	9.2
Blue runner (<i>Caranx crysos</i>)	2,863	1,164	2.5	2.5
Pink cusk-eel (<i>Genypterus blacodes</i>)	48,684	50,00-20,000	2.4	9.7
White hake (<i>Urophycis tenuis</i>)	8,620	2,361-3,656	2.4	3.7
Pacific halibut (<i>Hippoglossus stenolepis</i>)	41,587	13,640-18,180	2.3	3.0
Blackspot(=red) seabream (<i>Pagellus bogaraveo</i>)	1,790	658-772	2.3	2.7
Northern red snapper (<i>Lutjanus campechanus</i>)	4,503	2,000	2.3	2.3
Kingklip (<i>Genypterus capensis</i>)	9,965	3,300-4,500	2.2	3.0
Australian salmon (<i>Arripis trutta</i>)	7,347	200-3,500	2.1	37
Blotched picarel (<i>Spicara maena</i>)	635	19-298	2.1	34
Lumpfish(=Lumpsucker) (<i>Cyclopterus lumpus</i>)	14,979	2,000-7,000	2.1	7.5
Giant catfish (<i>Netuma thalassina</i>)	585	220-300	2.0	2.7
Common warehou (<i>Seriolella brama</i>)	4,373	2,200	2.0	2.0
European eel (<i>Anguilla anguilla</i>)	6,603	3,500	1.9	1.9
Swordfish (<i>Xiphias gladius</i>)	107,980	31,000-60,000	1.8	3.5
River lamprey (<i>Lampetra fluviatilis</i>)	144	48-81	1.8	3.0
Blue shark (<i>Prionace glauca</i>)	30,300	14,412-16,562	1.8	2.1

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Common dolphinfish (<i>Coryphaena hippurus</i>)	49,628	28,000	1.8	1.8
Gulf kingcroaker (<i>Menticirrhus littoralis</i>)	807	454	1.8	1.8
Red hake (<i>Urophycis chuss</i>)	2,652	1,500	1.8	1.8
Argentine goatfish (<i>Mullus argentinae</i>)	256	110-149	1.7	2.3
Giant stargazer (<i>Kathetostoma giganteum</i>)	3,458	2,000	1.7	1.7
Spotted weakfish (<i>Cynoscion nebulosus</i>)	3,403	2,000	1.7	1.7
Aba (<i>Gymnarchus niloticus</i>)	8,844	4,070-5,550	1.6	2.2
Rainbow runner (<i>Elagatis bipinnulata</i>)	15,983	10,000	1.6	1.6
Great barracuda (<i>Sphyræna barracuda</i>)	14,018	907-9,072	1.5	15.
Cape gurnard (<i>Chelidonichthys capensis</i>)	650	436	1.5	1.5
Rex sole (<i>Glyptocephalus zachirus</i>)	2,203	1,500	1.5	1.5
Blue ling (<i>Molva dypterygia</i>)	12,224	6,600-9,000	1.4	1.9
Bastard halibut (<i>Paralichthys olivaceus</i>)	8,896	6,042-6,298	1.4	1.5
Chinook(=Spring=King)salmon (<i>Oncorhynchus tshawytscha</i>)	11,529	8,341	1.4	1.4
Indo-Pacific tarpon (<i>Megalops cyprinoides</i>)	1,512	1,062-1,065	1.4	1.4
Gilthead seabream (<i>Sparus aurata</i>)	8,068	590-6,254	1.3	14
Barred grunt (<i>Conodon nobilis</i>)	180	141	1.3	1.3
King mackerel (<i>Scomberomorus cavalla</i>)	12,721	10,000	1.3	1.3
Lake(=Common)whitefish (<i>Coregonus clupeaformis</i>)	12,612	10,000	1.3	1.3
Squeteague(=Gray weakfish) (<i>Cynoscion regalis</i>)	1,456	1,134	1.3	1.3
Tuna-like fishes nei (<i>Scombroidei</i>)*	208,362	54-180,000	1.2	3,900
Bearded brotula (<i>Brotula barbata</i>)	1,616	909-1,364	1.2	1.8
Bobo mullet (<i>Joturus pichardi</i>)	296	247	1.2	1.2
Brushtooth lizardfish (<i>Saurida undosquamis</i>)	384	21-361	1.1	18
European conger (<i>Conger conger</i>)	15,622	1,718-14,854	1.1	9.1
Shallow-water Cape hake (<i>Merluccius capensis</i>)	2,260	490-2,030	1.1	4.6
Bluespot mullet (<i>Valamugil seheli</i>)	2,175	500-2,000	1.1	4.4
Bluestripe herring (<i>Herklotsichthys quadrimaculatus</i>)	27	13-25	1.1	2.2
Canary drum (=Baardman) (<i>Umbrina canariensis</i>)	2,550	2,400	1.1	1.1
Eastern Pacific bonito (<i>Sarda chiliensis chiliensis</i>)	4,538	4,000	1.1	1.1
European seabass (<i>Dicentrarchus labrax</i>)	10,225	300-10,000	1.0	34
Angler(=Monk) (<i>Lophius piscatorius</i>)	28,964	30,000	0.97	0.97

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Baird's slickhead (<i>Alepocephalus bairdii</i>)	2,254	33-2,431	0.93	69.
Atlantic salmon (<i>Salmo salar</i>)	3,917	4,500	0.87	0.87
Batfishes (<i>Platax spp</i>)*	2,919	775-3,521	0.83	3.8
Atlantic sabretooth anchovy (<i>Lycengraulis grossidens</i>)	50	19-61	0.82	2.6
Silver carp (<i>Hypophthalmichthys molitrix</i>)	10,523	12,794	0.82	0.82
West African Spanish mackerel (<i>Scomberomorus tritor</i>)	1,405	1,711	0.82	0.82
Brill (<i>Scophthalmus rhombus</i>)	2,800	3,475	0.81	0.81
Argentinian sandperch (<i>Pseudoperca semifasciata</i>)	2,414	2,200-3,000	0.80	1.1
Masu(=Cherry) salmon (<i>Oncorhynchus masou masou</i>)	1,570	2,000	0.79	0.79
Tench (<i>Tinca tinca</i>)	3,087	3,900	0.79	0.79
Lake cisco (<i>Coregonus artedi</i>)	646	832	0.78	0.78
Pacific jack mackerel (<i>Trachurus symmetricus</i>)	1,435	190-1,866	0.77	7.5
Indo-Pacific sailfish (<i>Istiophorus platypterus</i>)	20,954	13,640-27,270	0.77	1.5
Blackhead seabream (<i>Acanthopagrus schlegelii schlegelii</i>)	730	704-960	0.76	1.0
Rainbow trout (<i>Oncorhynchus mykiss</i>)	3,744	500-5,000	0.75	7.5
Widow rockfish (<i>Sebastes entomelas</i>)	1,287	300-1,814	0.71	4.3
Blackmouth catshark (<i>Galeus melastomus</i>)	197	65-280	0.71	3.0
Petrale sole (<i>Eopsetta jordani</i>)	2,055	1,040-3,000	0.69	2.0
American eel (<i>Anguilla rostrata</i>)	935	1,364	0.69	0.69
Black cardinal fish (<i>Epigonus telescopus</i>)	2,723	900-4,138	0.66	3.0
Grass carp(=White amur) (<i>Ctenopharyngodon idella</i>)	23,689	29,540-36,360	0.65	0.80
Turbot (<i>Psetta maxima</i>)	7,165	2,065-11,225	0.64	3.5
Sea trout (<i>Salmo trutta trutta</i>)	3,855	6,000	0.64	0.64
Red grouper (<i>Epinephelus morio</i>)	1,395	1,364-2,273	0.61	1.0
Common dentex (<i>Dentex dentex</i>)	1,387	1,260-2,350	0.59	1.1
Black seabass (<i>Centropristis striata</i>)	1,605	227-2,750	0.58	7.1
Chilipepper rockfish (<i>Sebastes goodei</i>)	264	339-462	0.57	0.78
Blue antimora (<i>Antimora rostrata</i>)	62	87-116	0.54	0.72
Southern bluefin tuna (<i>Thunnus maccoyii</i>)	15,054	7,000-28,333	0.53	2.2
Cachama (<i>Colossoma macropomum</i>)	4,552	8,836	0.52	0.52
Cobia (<i>Rachycentron canadum</i>)	9,128	6,364-18,182	0.50	1.4
Freshwater drum (<i>Aplodinotus grunniens</i>)	496	1,000	0.50	0.50
Bluenose warehou (<i>Hyperoglyphe antarctica</i>)	2,882	6,000	0.48	0.48

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Bighead carp (<i>Hypophthalmichthys nobilis</i>)	1,506	3,306	0.46	0.46
Ladyfish (<i>Elops saurus</i>)	915	2,000	0.46	0.46
Sheepshead (<i>Archosargus probatocephalus</i>)	1,024	2,268	0.45	0.45
Argentine angelshark (<i>Squatina argentina</i>)	3,980	9,017	0.44	0.44
Asp (<i>Aspius aspius</i>)	1,151	2,720	0.42	0.42
American shad (<i>Alosa sapidissima</i>)	890	907-2,272	0.39	0.98
White grouper (<i>Epinephelus aeneus</i>)	943	2,419	0.39	0.39
Blackfin tuna (<i>Thunnus atlanticus</i>)	2,626	3,182-7,000	0.38	0.83
Argentine conger (<i>Conger orbignianus</i>)	213	428-583	0.37	0.50
Thornback ray (<i>Raja clavata</i>)	1,536	4,536	0.34	0.34
Cape elephantfish (<i>Callorhynchus capensis</i>)	527	1,155-1,575	0.33	0.46
Anglerfishes nei (<i>Lophiidae</i>)*	9,346	413-30,000	0.31	23
Grayling (<i>Thymallus thymallus</i>)	246	300-800	0.31	0.82
Hapuku wreckfish (<i>Polyprion oxygeneios</i>)	1,654	6,000	0.28	0.28
Malabar blood snapper (<i>Lutjanus malabaricus</i>)	453	1,633	0.28	0.28
Antarctic rockcods, noties nei (<i>Nototheniidae</i>)*	2,742	21-10,000	0.27	130
Boarfishes nei (<i>Caproidae</i>)*	31	35-116	0.27	0.88
Stargazer (<i>Uranoscopus scaber</i>)	122	111-460	0.26	1.1
Blue catfish (<i>Ictalurus furcatus</i>)	717	3,000	0.24	0.24
Slender rainbow sardine (<i>Dussumieria elopsoides</i>)	14	5-62	0.23	2.9
Buffalofishes nei (<i>Ictiobus spp</i>)*	1,560	908-6,810	0.23	1.7
Lingcod (<i>Ophiodon elongatus</i>)	2,941	9,072-13,608	0.22	0.32
Boxfishes nei (<i>Ostraciidae</i>)*	133	93-637	0.21	1.4
Atlantic halibut (<i>Hippoglossus hippoglossus</i>)	4,325	21,000	0.21	0.21
Argentine menhaden (<i>Brevoortia pectinata</i>)	72	162-349	0.20	0.44
Atlantic goldeneye tilefish (<i>Caulolatilus chrysops</i>)	190	951	0.20	0.20
Black drum (<i>Pogonias cromis</i>)	2,759	13,636	0.20	0.20
Striped bass (<i>Morone saxatilis</i>)	3,093	6,820-15,900	0.19	0.45
Tope shark (<i>Galeorhinus galeus</i>)	4,812	25,000	0.19	0.19
Black catfishes nei (<i>Chrysichthys spp</i>)*	7,337	3-40,500	0.18	2,400
Great Northern tilefish (<i>Lopholatilus chamaeleonticeps</i>)	877	2,000-5,000	0.18	0.44
Blue marlin (<i>Makaira nigricans</i>)	33,134	180,000	0.18	0.18
Law croaker (<i>Pseudotolithus senegallus</i>)	1,077	6,080	0.18	0.18

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Alexandria pompano (<i>Alectis alexandrinus</i>)	488	2,851	0.17	0.17
Striped bonito (<i>Sarda orientalis</i>)	509	3,000	0.17	0.17
Bonefish (<i>Albula vulpes</i>)	517	3,333	0.16	0.16
Atlantic emperor (<i>Lethrinus atlanticus</i>)	78	509-511	0.15	0.15
Atlantic bluefin tuna (<i>Thunnus thynnus</i>)	35,935	262,000	0.14	0.14
Channel catfish (<i>Ictalurus punctatus</i>)	250	2,000	0.13	0.13
Atlantic tomcod (<i>Microgadus tomcod</i>)	32	108-258	0.12	0.30
Atlantic sailfish (<i>Istiophorus albicans</i>)	2,753	18,140-27,220	0.10	0.15
Two-spot red snapper (<i>Lutjanus bohar</i>)	282	2,759-2,795	0.10	0.10
Pacific bluefin tuna (<i>Thunnus orientalis</i>)	9,786	38,000-99,300	0.099	0.26
Greater amberjack (<i>Seriola dumerili</i>)	2,451	8,182-27,300	0.090	0.30
Antarctic toothfish (<i>Dissostichus mawsoni</i>)	2,312	28,000	0.083	0.083
Nassau grouper (<i>Epinephelus striatus</i>)	358	4,540	0.079	0.079
Wahoo (<i>Acanthocybium solandri</i>)	3,017	4,536-38,636	0.078	0.67
Black skipjack (<i>Euthynnus lineatus</i>)	550	695-7,165	0.077	0.79
Blueback shad (<i>Alosa aestivalis</i>)	5	44-60	0.076	0.10
Splendid alfonsino (<i>Beryx splendens</i>)	247	3,102-3,239	0.076	0.080
Antarctic silverfish (<i>Pleuragramma antarcticum</i>)	2	21	0.074	0.074
Black rockfish (<i>Sebastes melanops</i>)	103	1,056-1,440	0.071	0.097
Striped marlin (<i>Tetrapturus audax</i>)	7,845	56,750-113,500	0.069	0.14
Black marlin (<i>Makaira indica</i>)	5,861	85,000-90,000	0.065	0.069
Brown bullhead (<i>Ameiurus nebulosus</i>)	10	35-168	0.060	0.28
Dusky grouper (<i>Epinephelus marginatus</i>)	2,583	44,079	0.059	0.059
Broad-striped anchovy (<i>Anchoa hepsetus</i>)	1	16	0.056	0.056
Canary rockfish (<i>Sebastes pinniger</i>)	111	2,000	0.056	0.056
Brazilian groupers nei (<i>Mycteroperca</i> spp)*	1,433	660-26,000	0.055	2.20
Mountain mullet (<i>Agonostomus monticola</i>)	7	128	0.054	0.054
Yellowtail amberjack (<i>Seriola lalandi</i>)	790	4,397-15,000	0.053	0.18
Dusky smooth-hound (<i>Mustelus canis</i>)	332	6,752	0.049	0.049
Birdbeak dogfish (<i>Deania calcea</i>)	202	1,242-4,253	0.047	0.16
Wels(=Som)catfish (<i>Silurus glanis</i>)	9,153	100,000-200,000	0.046	0.092
Geelbek croaker (<i>Atractoscion aequidens</i>)	483	11,000	0.044	0.044
Wreckfish (<i>Polyprion americanus</i>)	923	22,600	0.041	0.041
California flounder (<i>Paralichthys californicus</i>)	399	7,260-9,900	0.040	0.055

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Shortfin mako (<i>Isurus oxyrinchus</i>)	4,694	60,000-135,000	0.035	0.078
Barbel (<i>Barbus barbus</i>)	124	1,800-3,636	0.034	0.069
Black dogfish (<i>Centroscyllium fabricii</i>)	78	1,085-2,328	0.033	0.072
Dogtooth tuna (<i>Gymnosarda unicolor</i>)	653	15,000-20,000	0.033	0.044
Black goby (<i>Gobius niger</i>)	2	63	0.032	0.032
Arctic char (<i>Salvelinus alpinus alpinus</i>)	135	454-4,545	0.030	0.30
Angelsharks, sand devils nei (<i>Squatinae</i>)*	532	1,684-19,794	0.027	0.32
Atlantic white marlin (<i>Tetrapturus albidus</i>)	731	27,215	0.027	0.027
Barbus cyclolepis (<i>Barbus cyclolepis</i>)	2	44-94	0.026	0.056
Blonde ray (<i>Raja brachyura</i>)	104	3,146-4,290	0.024	0.033
Tarpon (<i>Megalops atlanticus</i>)	952	40,000	0.024	0.024
Brown meagre (<i>Sciaena umbra</i>)	40	524-1,790	0.022	0.076
Cabezon (<i>Scorpaenichthys marmoratus</i>)	93	3,080-4,200	0.022	0.030
Leaping mullet (<i>Liza saliens</i>)	8	128-402	0.020	0.063
Blacktip shark (<i>Carcharhinus limbatus</i>)	294	18,000	0.016	0.016
Starry sturgeon (<i>Acipenser stellatus</i>)	147	8,953	0.016	0.016
Blacktip grouper (<i>Epinephelus fasciatus</i>)	3	138-167	0.015	0.019
Opah (<i>Lampris guttatus</i>)	706	50,000	0.014	0.014
Blue skate (<i>Dipturus batis</i>)	596	5,232-44,213	0.013	0.11
Cero (<i>Scomberomorus regalis</i>)	67	5,000	0.013	0.013
Antarctic starry skate (<i>Amblyraja georgiana</i>)	20	1,201-1,731	0.012	0.017
Bocaccio rockfish (<i>Sebastes paucispinis</i>)	34	2,119-2,889	0.012	0.016
Allis shad (<i>Alosa alosa</i>)	30	2,700	0.011	0.011
Porbeagle (<i>Lamna nasus</i>)	1,542	135,000	0.011	0.011
California sheephead (<i>Semicossyphus pulcher</i>)	50	3,520-4,800	0.010	0.014
Yellowedge grouper (<i>Epinephelus flavolimbatus</i>)	40	2,270-4,540	0.0088	0.018
Biglip grunt (<i>Plectorhinchus macrolepis</i>)	4	472	0.0087	0.0087
White sturgeon (<i>Acipenser transmontanus</i>)	157	9,072-18,144	0.0086	0.017
Bar jack (<i>Carangoides ruber</i>)	7	454-909	0.0082	0.016
Smooth-hound (<i>Mustelus mustelus</i>)	135	4,508-19,539	0.0069	0.030
Bullet tuna (<i>Auxis rochei rochei</i>)	6	861-890	0.0062	0.0065
American conger (<i>Conger oceanicus</i>)	44	4,545-9,090	0.0049	0.0098
Thresher (<i>Alopias vulpinus</i>)	473	113,500	0.0042	0.0042
Bigspined boarfish (<i>Pentaceros decacanthus</i>)	4	500-1,000	0.0041	0.0082

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Blackbellied angler (<i>Lophius budegassa</i>)	9	1,631-2,120	0.0040	0.0052
Bigeye grenadier (<i>Macrourus holotrachys</i>)	2	468	0.0040	0.0040
Flat needlefish (<i>Ablennes hians</i>)	9	2,036-2,900	0.0033	0.0046
Arapaima (<i>Arapaima gigas</i>)	239	73,100	0.0033	0.0033
Scalloped hammerhead (<i>Sphyrna lewini</i>)	365	115,102	0.0032	0.0032
Oceanic whitetip shark (<i>Carcharhinus longimanus</i>)	212	35,000-70,000	0.0030	0.0060
Broomtail grouper (<i>Mycteroperca xenarcha</i>)	79	20,020-27,300	0.0029	0.0040
Ballan wrasse (<i>Labrus bergylta</i>)	5	1,000-2,000	0.0024	0.0049
Red drum (<i>Sciaenops ocellatus</i>)	7	1,818-3,636	0.0020	0.0039
Blackfin icefish (<i>Chaenocephalus aceratus</i>)	2	788-929	0.0018	0.0021
Black carp (<i>Mylopharyngodon piceus</i>)	27	15,000	0.0018	0.0018
Brook trout (<i>Salvelinus fontinalis</i>)	8	500-5,000	0.0016	0.016
Black driftfish (<i>Hyperoglyphe bythites</i>)	4	1,364-2,727	0.0016	0.0032
Angelshark (<i>Squatina squatina</i>)	22	14,000-18,000	0.0012	0.0016
Snowy grouper (<i>Epinephelus niveatus</i>)	5	3,636-4,545	0.0012	0.0015
Golden grey mullet (<i>Liza aurata</i>)	1	924	0.0012	0.0012
Brown smooth-hound (<i>Mustelus henlei</i>)	3	471-2,685	0.0011	0.0064
Nurse shark (<i>Ginglymostoma cirratum</i>)	159	90,000-150,000	0.0011	0.0018
Sickle pomfret (<i>Taractichthys steindachneri</i>)	9	7,128-8,036	0.0011	0.0012
Gag (<i>Mycteroperca microlepis</i>)	9	909-9,091	0.0010	0.010
Bigeye thresher (<i>Alopias superciliosus</i>)	160	160,000	0.0010	0.0010
Beluga (<i>Huso huso</i>)	75	35,000-100,000	0.00075	0.0022
Warsaw grouper (<i>Epinephelus nigritus</i>)	26	13,636-36,364	0.00072	0.0019
Bathyraja rays nei (<i>Bathyraja</i> spp)*	2	277-3,180	0.00070	0.0080
Sandbar shark (<i>Carcharhinus plumbeus</i>)	50	60,000-96,000	0.00052	0.00083
Arctic skate (<i>Amblyraja hyperborea</i>)	1	1,201-1,731	0.00045	0.00065
Giant guitarfish (<i>Rhynchobatus djiddensis</i>)	85	190,000	0.00045	0.00045
Basketwork eel (<i>Diastobranchus capensis</i>)	1	2,250-3,068	0.00036	0.00049
Broadnose sevengill shark (<i>Notorynchus cepedianus</i>)	4	22,613	0.00019	0.00019
Yellowfin grouper (<i>Mycteroperca venenosa</i>)	1	6,810	0.00013	0.00013
Black grouper (<i>Mycteroperca bonaci</i>)	3	26,000	0.00012	0.00012
Basking shark (<i>Cetorhinus maximus</i>)	246	3,000,000	0.000082	0.000082
King of herrings (<i>Regalecus glesne</i>)	6	59,840-81,600	0.000075	0.00010
Tiger shark (<i>Galeocerdo cuvier</i>)	38	385,000-685,000	0.000055	0.000099

Table 4. Numbers of fish caught estimated from FAO capture tonnages and estimated mean weights (cont.)

Species (scientific name)	Average annual capture 1999-2007 (t) (FAO, 2009a)	Estimated mean weight range (g)	Estimated number range (lower) in millions	Estimated number range (upper) in millions
Bluntnose sixgill shark (<i>Hexanchus griseus</i>)	9	500,000	0.000018	0.000018
Ocean sunfish (<i>Mola mola</i>)	11	1,000,000	0.000011	0.000011
Great white shark (<i>Carcharodon carcharias</i>)	2	680,000	0.0000025	0.0000025
Total for species with an estimated mean weight	53,025,428		680,000	2,000,000
Total for species without an estimated mean weight	24,362,894		300,000	760,000
Total fish capture	77,388,322		970,000	2,700,000